


PHYSICS PAPER 232/1 K.C.S.E. 2002 MARKING SCHEME

1. 11.72/11.72 CM/0.01172M
2.




3. g moves / shifts to the right / C.O.M. moves/ shifts/ more weight or mass of he right/ weight will have a clockwise movement about O/causing greater moment of force towards right than left.

4. $R = V = 0.35 = 0.5\Omega$
 $I = 0.70$
 $P = \frac{RA}{C} = \frac{0.5 \times 8 \times 10^{-3}}{0.5} = 8 \times 10^{-3} \Omega \text{ m.}$

5. $p = \frac{F}{A} = \frac{2500}{0.025} = 100,000 \text{ Pa}$
 $P = \frac{F}{A} = \frac{2500}{0.025} = 100,000 \text{ Pa}$
 Total press = $\frac{2500}{0.025} = 100,000 \text{ Pa} = 100,000 \text{ N/m}^2$

6. -Low temperature reduces K.E / velocity of molecules
 - Hence lower rate of collision / less collision -Reduction in pressure
7. Can B Good absorber of radiation.

8.



A' & B' are the refracted rays

19. (Assume no heat losses)


Heat gained = heat lost
 $2 \times c \times (30 - 20) = 90 \times 15 \times 60$
 $C = \frac{90 \times 15 \times 60}{20}$

$C = 4050 \text{ J/kgK}$

$E = pt = mc\Delta\theta$
 $90 \times 15 \times 60 = 2 \times c \times 10$
 $4050 \text{ J/kgK} = c$

20. Mattress increases stopping time/time of collision increased this reduces the rate of change of momentum.


21. $C = C_1 + C_2$ $Q = CV$
 $CT = 3 \times 2$ $5\mu F$ $V = \frac{Q}{C}$ $V = 1 \times 10^{-4} = 20V$



24. $V = f\lambda$
 $\lambda = \frac{v}{f} = \frac{330}{30} = 11m$

25. Law of floatation – a floating body displaces its own weight
 Weight of block = weight of mercury displaced
 $0.250 \times g = 13.6g$
 $\frac{0.25}{13.6 \times 10^3} = v$

$V = \frac{1.838 \times 10^{-5} m^3}{1.839 \times 10^{-5} m^3} = 18.4cm^3$



29. $p = VI$

	Kettle	Iron box	TV
$I = p/n = \frac{2500}{250} = 8A$		$750/250 = 3A$	$300/250 = 1.2A$
Total = 8 + 3 + 1.2 = 12.2A		= Appropriate fuse = 15A	

31. $107 - 42 = 65$

32. Penetrating power
33. Downwards
34. Work function of metal / min energy required to eject e-1 for excess energy work function.

[HTTP://ATIKASCHOOL.ORG](http://ATIKASCHOOL.ORG) - 0728 450 425

PHYSICS PAPER 232/2 K.C.S.E 2002 MARKING SCHEME

1a) (speed of light in vacuum $c = 3.0 \times 10^8 \text{ ms}^{-1}$)
 Refractive index = speed of light in vacuum
 $= \frac{3.0 \times 10^8 \text{ m/s}}{1.88 \times 10^8 \text{ m/s}}$
 $= 1.596 = 1.60$

b) $\sin C = \frac{1}{n}$
 $C = \sin^{-1} \left(\frac{1}{1.596} \right)$
 $C = 38.8^\circ - 38.48^\circ$
 $38.7 - 38.42$


c) $\frac{\sin \theta}{\sin 21.1} = 1.596$
 $\frac{\sin \theta}{\sin 21.1} = n$
 $\sin \theta = 1.596 \times \sin 21.1$
 $\theta = \sin^{-1} (0.58) = 35.25^\circ - 35.15^\circ$
 $35.35^\circ - 35.21^\circ$

2. β - beta radiation
 Force is of the circle implying negatively charged (Fleming's left hand rule)

(bi) K= alpha (ii) X= 88 Y= 288

- (ci) Increase in thickness
- (ii) Increase in thickness reduces the radiation reaching the Geiger tube
- (iii) Increase in pressure
- (iv) Increase roller pressure squeezes metal sheet (possess more) reducing the thickness of foil coming out of them.
- (v) Alpha particles have little penetration very few or none pass though foil.
- (vi)

3.



a i) R- to pass through the c.o.g Forces not labeled. Award half for each
 (ii) $= mg \sin \theta = 30.0 \times 10 \sin 10^\circ = 52.1 \text{ N}$ (accept 52.08, 52.08, 52.09)

(ii) $A = F$ Net force down = $Mg \sin \theta - \text{friction} = 52.1 - 20 = 32.1$

$M = \frac{32.1}{3.0} = 1.07 \text{ M/S}^2$

(iii) Acceleration increases with the increase in angle

- 4 a i) A ice absorbs latent heat without in temperature (or ice melting no change of temperature heat goes to latent heat fusion)
B Water molecules gain K.E (increase in K.E.)
C heat is used to change water into vapour.

ii) Water has anomalous expansion, where we have maximum density at 4°C.
Anomalous behaviour/explain.

iii) Frozen seawater has a lower temperature than frozen fresh water boiling point of sea water is higher than fresh water.

(b) (heat gained = ML + MCθ
= 3 x 336 x 10³ + 3 x 4200x5
= 1.07 x 10⁶J

5 a i) Transverse waves (accept elliptical)

ii) As waves move in the medium, the particles of medium do not move: they vibrate in positions so cork does not move.

iii) Period of wave T= 0.205

$$f = \frac{1}{T} = 5\text{Hz}$$

$$V = f\lambda$$

$$\lambda = \frac{0.30}{5} = 0.60\text{M}$$

iv) Velocity decreases when depth decreases hence the λ decreases (since frequency is constant wavelength decreases)

b) 1st resonance $\frac{\lambda}{4} = I_2 - I_1$ $\lambda = \frac{2(I_2 - I_1)}{1}$ OR $V = 2F(I_2 - I_1)$
f = $\frac{V}{2(I_2 - I_1)}$
 $\frac{\lambda}{2} = 129 - 77$

2nd resonance $3\lambda = I_2 + C$ $\lambda = 104 \text{ cm}$ =340
 $V = f\lambda$
 $340 = f \times 1.04 = 326.9 \text{ Hz.}$
 $F = 327 \text{ Hz (326.9)}$

a) Charles law: for a fixed mass of a gas at a constant pressure the volume is directly proportional to the absolute temperature Kelvin thermodynamics.

bi) Volume of gas trapped by drop of cone sulphuric acid, water in heated (in both) and volume (height) of gas: in tube increase as temperature rises; values of height H and T are tabulated; a graph of volume V versus temperature T°C is plotted;

graph is straight line cutting T at -273°C (absolute Zero); so volume is directly proportional to absolute temperature.

ii) -Short temperature range - Keeping pressure constant is difficult

ci) When $\theta - \theta T - 273\text{k}$ Extrapolation on graph show:

Pressure read off $\beta = 9.7 \times 10^4 \text{ pa}$

ii) $p_1 = 1.15 \times 10^5 \text{ pa}$ $\theta_1 = 52.0^{\circ}\text{C}$

$p_2 = 1.25 \times 10^5 \text{ pa}$ $\theta_2 = 80.0^{\circ}\text{C}$

p_1	p_2
$T_0 + \theta_1$	$T_0 + \theta_2$
1.115×10^5	1.25×10^5
$T_0 + 52$	$T_0 + 80.0$
T_0 270	

- Rise in volume height

- Rise in temperature

-Recording of tabulation

- Graph

-Analysis of graph

-Conclusion

Alternatives

$$P = mx + c$$

$$P = k\theta + kT_0 \text{ when } K \text{ gradient.}$$

$$K = \frac{Dv}{Dx} = \frac{(1.14 - 1) \times 10^5}{50 - 10}$$

$$= \frac{0.14 \times 10^5}{40}$$

$$= \frac{14000}{40} = 350 \text{ pac ()}$$

$$KT = \text{Constant}$$

$$C = 9.6 \times 10^4$$

$$350 T_0 = 9.67 \times 10^4$$

$$T_0 = 274.3 \text{ (266-284)}$$

5. ai) μV light removes electrons on zinc plate. This lowers the excess charge constant (negative) on leaf leading to collapse/ becomes less negative (more positive)

ii) Since μv light removes electrons positive charge re attracts the electrons thus keeps the charge constant and so leaf does not collapse.

bi) Frequency of incident light / energy of photon / energy of light work function of surface

ii) From $K_{\text{max}} = hf - \theta$

h is slope of graph

$$\text{Slope} = (10 - 20) \times 10^{-19}$$

$$(2.6 - 1.4) \times 10^{15}$$

$$H = 6.7 \times 10^{-34} \text{ fs}$$

At $K_{\text{max}} = \theta$ $hf = 0$

Extrapolation shown or

$$\text{Read off } f_0 = 1.07 \times 10^{15} \text{ Hz}$$

$$\Theta = 1.07 \times 10^{15} \times 6.67 \times 10^{-34}$$

$$= 7.4 \times 10^{-19}$$

$$\begin{aligned}
 \text{c) } K_{\text{max}} &= hf - \theta \\
 &= \frac{6.67 \times 10^{-34} \times 5.5 \times 10^{14}}{1.6 \times 10^{-19}} \\
 &= 2.29 \text{ eV} \\
 &\text{Since } hf < \theta \text{ no photo electric effect} \\
 \text{Or } E &= hf = 6.67 \times 10^{-34} \times 5.5 \times 10^{14} \\
 \theta &= 2.5 \times 1.6 \times 10^{-19}
 \end{aligned}$$

[HTTP://ATIKASCHOOL.ORG](http://ATIKASCHOOL.ORG) - 0728 450 425