


5.0 THE YEAR 2012 KCSE EXAMINATION MARKING SCHEMES

5.1 MATHEMATICS (121 AND 122)



5.1.1 Mathematics Alternative A Paper 1 (121/1)

1.	$\frac{\frac{6}{5} - \frac{4}{3}}{\frac{1}{8} - \frac{1}{4}} - \frac{14}{15}$ $= \frac{-\frac{2}{15}}{-\frac{1}{8}} - \frac{14}{15}$ $= \frac{16}{15} - \frac{14}{15}$ $= \frac{2}{15}$	M1 M1 M1 A1 4	numerator denominator
2.	$\frac{1}{0.216} = 4.630$ $\frac{\sqrt[3]{0.512}}{0.216} = 0.8 \times 4.630$ $= 3.704$	B1 M1 A1 3	
3.	$(2x^2 - 3y^3)^2 + 12x^2y^3$ $= 4x^4 - 12x^2y^3 + 9y^6 + 12x^2y^3$ $= 4x^4 + 9y^6$	M1 A1 2	
4.	$\frac{24}{2} = \frac{1}{2} \times 8 \times x \sin 30^\circ$ $x = \frac{12}{4 \sin 30} = 6cm$ $perimeter = 2(6 + 8) = 28$	M1 M1 A1 3	or equivalent
5.	$9^{2y} \times 2^x = 9 \times 8$ $(3^2)^{2y} \times 2^x = 3^2 \times 2^3$ $(3^2)^{2y} = 3 \text{ and } 2^x = 2^3$ $4y = 2 \text{ and } x = 3$ $y = \frac{1}{2} \text{ and } x = 3$	M1 M1 A1 3	equating indices

6.	<p>LCM of 9, 15 and 21</p> $3^2 \times 5 \times 7 = 315 \text{ minutes}$ <p>Last time of ringing together</p> $\begin{array}{r} 11:00 \\ - 5:15 \\ \hline 5:45 \text{ p.m.} \end{array}$	<p>B1</p> <p>M1</p> <p>A1</p> <p>3</p>	<p>For 315 minutes</p> <p>For subtraction</p>
7.	$\frac{x}{8} = \frac{x}{20} + \frac{1}{4}$ $\frac{x}{8} - \frac{x}{20} = \frac{1}{4}$ $\Rightarrow \frac{3x}{40} = \frac{1}{4}$ $x = 3\frac{1}{3}$ <p>Distance to shopping centre</p> $12 - 3\frac{1}{3} = 8\frac{2}{3} \text{ km}$	<p>M1</p> <p>A1</p> <p>B1</p> <p>3</p>	
8.	 <p>Construction of <math>135^\circ</math> angle between lines <math>AB = 4 \text{ cm}</math> and <math>BC = 6 \text{ cm}</math></p> <p>Construction of <math>60^\circ</math> angle between lines <math>AB = 4 \text{ cm}</math> and <math>AD = 3 \text{ cm}</math></p> <p>Completion of quadrilateral ABCD</p> $\angle BCD = 31^\circ \pm 1^\circ$	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>4</p>	

9.	$\begin{pmatrix} -3 \\ -2 \end{pmatrix} - \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ $= \begin{pmatrix} -1 \\ -5 \end{pmatrix}$ <p>magnitude = <math>\sqrt{1^2 + (-5)^2}</math></p> $= \sqrt{26} \approx 5.1$	M1  M1 A1  3	
10.	$x = \tan^{-1} \frac{3}{7} = 23.20^\circ$ $\cos(90 - 23.2)^\circ = 0.3939$	B1  B1  2	
11.	$A^2 = \begin{pmatrix} 1 & 0 \\ -2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -2 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -8 & 9 \end{pmatrix}$ $2AB = 2 \begin{pmatrix} 1 & 0 \\ -2 & 3 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 2 & 1 \end{pmatrix} = 2 \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix} = \begin{pmatrix} 6 & 0 \\ 0 & 6 \end{pmatrix}$ $C = 2AB - A^2 = \begin{pmatrix} 6 & 0 \\ 0 & 6 \end{pmatrix} - \begin{pmatrix} 1 & 0 \\ -8 & 9 \end{pmatrix}$ $= \begin{pmatrix} 5 & 0 \\ 8 & -3 \end{pmatrix}$	B1  B1  M1  A1  4	
12.	$\log_{10} \left( \frac{x^2}{2^3} \times 32 \right) = 2$ $\frac{x^2}{2^3} \times 2^5 = 100$ $4x^2 = 100$ $x = \sqrt{25} = \pm 5$ $x = 5$	M1  M1  A1  3	dropping logs.

13.	$2y = 4x + 5 \Rightarrow y = 2x + \frac{5}{2}$ <p>gradient, <math>M_1</math> of line = 2</p> <p>gradient, <math>M_2</math>, of perpendicular is given by</p> $2M_2 = -1 \Rightarrow M_2 = -\frac{1}{2}$ <p>equation of line L</p> $\frac{y-1}{x-3} = -\frac{1}{2}$ $y = -\frac{1}{2}x + \frac{5}{2}$	B1 M1 A1	
14. (a) 195250 Chinese Yuan into Kenya Shillings $= 195250 \times 12.34 = 2409385$ (b) Balance: $= 2409385 - 1258000$ $= 1151385$ Balance in S.A. Rand $= \frac{1151385}{11.37}$ $= 101265$		B1 M1 M1 A1	
		3	
		4	

15.	<p>Volume of solid</p> $= \frac{1}{3} \times \frac{22}{7} \times 10.5^2 \times 15 - \frac{22}{7} \times 3.5^2 \times 8$ $= 1732.5 - 308$ $= 1424.5 \text{ cm}^3$	<p>M1 M1</p> <p>A1</p> <p>3</p>	
16.	$\left. \begin{aligned} 4(A - 2) &= B + 2 \\ 2(A + 10) &= B + 10 \end{aligned} \right\}$ $4A - B = 10 \dots (i)$ $\mp 2A \pm B = \pm 10 \dots (ii)$ <hr/> $2A = 20$ $\Rightarrow A = 10$ <p>Substitute A = 10 in (i)</p> $4 \times 10 - B = 10$ $\Rightarrow B = 30$	<p>M1</p> <p>M1</p> <p>A1</p> <p>3</p>	<p>for both values of A and B</p>
17. (a)	<p>modal class 40 - 44</p> <p>(b) (i) mid points:</p> <p>22, 27, 32, 37, 42, 47, 52, 57</p> $\frac{22 \times 2 + 27 \times 15 + 32 \times 18 + 37 \times 25 + 42 \times 30 + 47 \times 6 + 52 \times 3 + 57 \times 2}{101}$ $= 37.25$	<p>B1</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p>	<p>fx</p> <p>for <math>\frac{\Sigma fx}{\Sigma f}</math></p>


	<p>(ii) Cumulative frequencies</p> <p>2, 17, 35, 60, 90, 96, 99, 101</p> $\frac{16}{25} \times 5$ $= 3.2$ $34.5 + 3.2$ $= 37.7$ <p>difference <math>37.7 - 37.25</math></p> $= 0.45$	<p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>B1</p>	
		10	
18. (a)	$ AB  = \sqrt{169 - 25} = 12$	B1	
(b)	$2 \times 5 \times 12 + 2 \times 5 \times 15 + 2 \times 12 \times 15$ $= 630 \text{ cm}^2$	<p>M1</p> <p>M1</p> <p>A1</p>	3 pairs of congruent faces summing up
(c)	<p>volume = <math>5 \times 12 \times 15 \text{ cm}^3</math></p> <p>mass = <math>7.6 \times 5 \times 12 \times 15</math></p> $= 6840 \text{ gm}$ $= \frac{6840}{1000}$ $= 6.84 \text{ kg}$	<p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p>	division by 1000
(d)	$\frac{150 \times 120 \times 100 \text{ cm}^3}{15 \times 12 \times 5 \text{ cm}^3}$ $= 2000$	<p>M1</p> <p>A1</p>	
		10	

19. (a)	<p><i>Ratio: copper: zinc: tin</i></p> <table border="0"> <thead> <tr> <th>copper</th> <th>zinc</th> <th>tin</th> </tr> </thead> <tbody> <tr> <td>3</td> <td><math>\frac{2}{3}</math></td> <td>5</td> </tr> <tr> <td>9</td> <td>6</td> <td>10</td> </tr> </tbody> </table> <p>Copper : zinc : tin = 9 : 6 : 10</p>	copper	zinc	tin	3	$\frac{2}{3}$	5	9	6	10	M1 A1	
copper	zinc	tin										
3	$\frac{2}{3}$	5										
9	6	10										
(b) (i)	<p>mass of tin</p> $= 250 \times \frac{10}{25}$ $= 100\text{kg}$	M1 A1										
(ii)	<p>mass of zinc and tin in alloy B:</p> $\text{mass of copper} = \frac{70}{100} \times 90$ $= 63$ <p><math>\therefore</math> mass of zinc and tin:</p> $= 250 - 63$ $= 187$	M1 M1 A1										
(c)	<p>amount of tin in alloy A than B:</p> <p>mass of tin in alloy B</p> $= \frac{8}{11} \times 187$ $= 136$ <p>difference:</p> $136 - 100$ $= 36$	M1 A1										
		10										

20. (a)	$\frac{1}{x-2} - \frac{2}{x+5} = \frac{3}{x+1}$ $\frac{x+5-2(x-2)}{(x-2)(x+5)} = \frac{3}{x+1}$ $\frac{-x+9}{x^2+3x-10} = \frac{3}{x+1}$ $4x^2+x-39=0$ $(4x+13)(x-3)=0$ $x=3 \text{ or } x=-3\frac{1}{4}$	M1 A1 M1 A1	
(b)	<p>mean for second set of tests</p> $= \frac{147}{y+2}$ $\frac{120}{y} - \frac{147}{y+2} = 3$ $\frac{120y+240-147y}{y(y+2)} = 3$ $-27y+240=3y^2+6y$ $-9y+80=y^2+2y$ $y^2+11y-80=0$ $(y-5)(y+16)=0$ $y=5 \text{ or } -16$ <p>No. of tests: <math>5+2=7</math></p>	B1 M1 M1 A1 M1 A1	elimination of denominator     factorization
		A1	
		10	



21.



a) (i)  $OPQR$  ✓ drawn

B1

$O'P'Q'R'$  ✓ drawn

B1

(ii) Perpendicular bisectors ✓ drawn (at least 2)

B1

centre of rotation  $(0, -1)$  shown

B1

angle of rotation  $-90^\circ$

B1

b) line of reflection  $x = y$  drawn

B1

can be implied

quadrilateral  $O'P'Q'R'$  drawn

B1

c) (i) directly congruent quads:

$OPQR$  and  $O'P'Q'R'$

B1

(ii) Oppositely congruent quads.:

$OPQR$  and  $O''P''Q''R''$

B1


$O'P'Q'R'$  and  $O''P''Q''R''$

B1

10

22. (a) (i)	<p>x - intercepts</p> <p>when <math>y=0</math></p> $x^2(2x + 3) = 0$ $x = 0 \text{ and } x = -\frac{3}{2}$	M1																	
(ii)	<p>y - intercept</p> <p>when <math>x = 0, y = 0</math></p>	A1																	
(b) (i)	<p>stationary points of curve</p> $\frac{dy}{dx} = 6x^2 + 6x$ <p>stationery points when <math>\frac{dy}{dx} = 0</math></p> <p>i.e. <math>6x^2 + 6x = 0</math></p> $6x(x + 1) = 0$ <p><math>x = 0</math> or <math>x = -1</math></p> <p><math>\therefore</math> stationary points are:</p> <p><math>(0,0)</math> and <math>(-1,1)</math></p>	M1																	
(ii)	<table border="1" data-bbox="328 1245 959 1413"> <tr> <td><math>x</math></td> <td>-2</td> <td><math>-1\frac{1}{2}</math></td> <td>-1</td> <td><math>-\frac{1}{2}</math></td> <td>0</td> <td><math>\frac{1}{2}</math></td> <td>1</td> </tr> <tr> <td><math>\frac{dy}{dx}</math></td> <td>12</td> <td><math>4\frac{1}{2}</math></td> <td>0</td> <td><math>-1\frac{1}{2}</math></td> <td>0</td> <td><math>4\frac{1}{2}</math></td> <td>12</td> </tr> </table> <p>minimum point <math>(0,0)</math> maximum point <math>(-1,1)</math></p>	$x$	-2	$-1\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{dy}{dx}$	12	$4\frac{1}{2}$	0	$-1\frac{1}{2}$	0	$4\frac{1}{2}$	12	A1	checking points
$x$	-2	$-1\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1												
$\frac{dy}{dx}$	12	$4\frac{1}{2}$	0	$-1\frac{1}{2}$	0	$4\frac{1}{2}$	12												
		B1	for both																

(c)



points plotted at  $(-1\frac{1}{2}, 0)$ ,  $(-1, 1)$  and  $(0, 0)$


smooth curve

B1

B1

10

23. (a)



- ✓ location of R
- ✓ location of T
- complete  $\Delta$

(b) (i) Distance TS:  $6.6(\pm 1) \text{ cm}$   
 conversion  $6.6 \times 60 = 396 \text{ m}$

(ii) Bearing of T from S  
 $180 + 41^\circ(\pm 1^\circ) = 221^\circ$

(c) area of field  
 $\angle TRS = 60^\circ$   

$$\text{area} = \frac{1}{2} \times 300 \times 450 \sin 60^\circ$$

$$= \frac{58456.71476}{10000}$$

$$= 5.8 \text{ ha}$$

B1 length 5 cm and bearing  $300^\circ$

B1 length 7.5 cm; south of R

B1

B1

B1

B1

B1

M1

M1

A1

10

24. (a)	<p>length of RT:</p> $= \frac{3}{5} \times 10$ $= 6 \text{ cm}$	M1	
		A1	
	(b) (i) Perpendicular distance between PQ & RS		
	$= 10 \sin 40$	M1	
	$= 6.4 \text{ cm}$	A1	
	(ii) $\frac{TS}{\sin 40} = \frac{6}{\sin 60}$		
	$TS = \frac{6 \times \sin 40}{\sin 60}$	M1	
	$= 4.5 \text{ cm}$	A1	
	(c) length RS using cosine rule		
	$RS^2 = 6^2 + 4.5^2 - 2 \times 6 \times 4.5 \cos 80$	M1	
	$= 46.87299841$		
	$RS = 6.8$	A1	
	(d) area of $\Delta RST$		
	$= \frac{1}{2} \times 6 \times 4.5 \sin 80$	M1	
	$= 13.3$	A1	
		10	